arXiv:1403.2151v2 [gr-qc] 11 Mar 2014

Thermodynamics of phase transition in higher dimensional Reissner-Nordström-de Sitter black hole

Li-Chun Zhang, Meng-Sen Ma, Hui-Hua Zhao, Ren Zhao*

Institute of Theoretical Physics, Shanxi Datong University, Datong 037009, China and

Department of Physics, Shanxi Datong University, Datong 037009, China

It is well known that there are black hole and the cosmological horizons for the Reissner-Nordström-de Sitter spacetime. Although the thermodynamic quantities on the horizons are not irrelevant, they satisfy the laws of black hole thermodynamics respectively. In this paper by considering the relations between the two horizons we give the effective thermodynamic quantities in $(n + 2)$-dimensional Reissner-Nordström-de Sitter spacetime. The thermodynamic properties of these effective quantities are analyzed, moreover, the critical temperature, critical pressure and critical volume are obtained. We carry out an analytical check of Ehrenfest equations and prove that both Ehrenfest equations are satisfied. So the spacetime undergoes a second order phase transition at the critical point. This result is consistent with the nature of liquid–gas phase transition at the critical point, hence deepening the understanding of the analogy of charged dS spacetime and liquid–gas systems.

Keywords: Reissner-Nordström-de Sitter black hole, the effective thermodynamic quantities, critical phenomena, stability

PACS numbers: 04.70.-s, 05.70.Ce

I. INTRODUCTION

Black hole physics, specially black hole thermodynamics, refer to the theories of gravity, statistical mechanics, particle physics and field theory and so on. This makes it receive a lot of attention[1–6]. Although the statistical explanation of the thermodynamic states of black holes is still lack, the properties of black hole thermodynamics are worth studying deeply,

* Email: zhao2969@sina.com

such as Hawking-Page phase transition[7], the critical phenomena of black holes. More interestingly, recently through the study of RN-AdS black hole it is shown that there exists similar phase transition to the one in the van der Waals-Maxwell vapor-liquid system[8, 9].

Motivated by the AdS/CFT correspondence[10], where the transitions have been related with the holographic superconductivity[11, 12], the subject that the phase transitions of black holes in asymptotically anti de-Sitter (AdS) spacetime has received considerable attention[13–17]. The underlying microscopic statistical interaction of the black holes is also expected to be understood via the study of the gauge theory living on the boundary in the gauge/gravity duality.

Recently, by considering the cosmological constant correspond to pressure in general thermodynamic system, namely

$$P = -\frac{1}{8\pi}\Lambda = \frac{3}{8\pi}\frac{1}{l^2}, \tag{1.1}$$

the thermodynamic volumes in AdS and dS spacetime are obtained[18–24]. The studies on phase transition of black holes have aroused great interest[16, 25–42]. Connecting the thermodynamic quantities of AdS black holes to ($P \sim V$)in the ordinary thermodynamic system, the critical behaviors of black holes can be analyzed and the phase diagram like van der Waals vapor-liquid system can be obtained. This helps to further understand the black hole entropy, temperature, heat capacities, etc. It also has a very important significance in completing the geometric theory of black hole thermodynamics.

As is well known, there are black hole horizon and the cosmological horizon in the appropriate range of parameters for de Sitter spacetime. The both horizons have thermal radiation, but with different temperatures. The thermodynamic quantities on the both horizons satisfy the first law of thermodynamics, and the corresponding entropy fulfill the area formula[22, 43, 44]. In recent years, the researches on the thermodynamic properties of de Sitter spacetime have drawn many attentions[16, 22, 43–48]. In the inflation epoch of early universe, the universe is a quasi-de Sitter spacetime. The cosmological constant introduced in de Sitter space may come from the vacuum energy, which is also a kind of energy. If the cosmological constant is the dark energy, the universe will evolve to a new de Sitter phase. To depict the whole history of evolution of the universe, we should have some knowledge on the classical and quantum properties of de Sitter space[22, 44, 49, 50].

Firstly, we expect the thermodynamic entropy to satisfy the Nernst theorem[45, 47, 51].

At present a satisfactory explanation to the problem that the thermodynamic entropy of the horizon of the extreme de Sitter spacetime do not fulfill the Nernst theorem is still lacked. Secondly, when considering the correlation between the black hole horizon and the cosmological horizon whether the thermodynamic quantities in de Sitter spacetime still have the phase transition and critical behavior like in AdS black holes. Thus it is worthy of our deep investigation and reflection to establish a consistent thermodynamics in de Sitter spacetime.

Because the thermodynamic quantities on the black hole horizon and the cosmological one in de Sitter spacetime are the functions of mass M, electric charge Q and cosmological constant Λ. The quantities are not independent each other. Considering the relation between the thermodynamic quantities on the two horizons is very important for studying the thermodynamic properties of de Sitter spacetime. Based on the relation we give the effective temperature and pressure of $(n+2)$-dimensional Reissner-Nordström-de Sitter(RN-dS) spacetime and analyze the critical behavior of the effective thermodynamic quantities.

The paper is arranged as follows: in the Sec.II we introduce the $(n+2)$-dimensional Reissner-Nordströmde Sitter(RN-dS) spacetime, and give the two horizons and corresponding thermodynamic quantities; In Sec.III by considering the relations between the two horizons we obtain the effective temperature and the equivalent pressure; In Sec. IV the critical phenomena of effective thermodynamic quantities is discussed; To investigate the nature of the phase transition at the critical point, we will introduce the classical Ehrenfest scheme and carry out an analytical check of both equations in Section V. Finally we discuss and summarize our results in Sec.VI. (we use the units $G_{n+1} = \hbar = k_B = c = 1$)

II. RN-DS SPACETIME

The line element of the $(n+2)$ dimensional RN-dS black holes is given by[43]

$$ds^2 = -f(r)dt^2 + f^{-1}dr^2 + r^2 d\Omega_n^2, \tag{2.1}$$

where

$$\begin{aligned} f(r) &= 1 - \frac{\omega_n M}{r^{n-1}} + \frac{n\omega_n^2 Q^2}{8(n-1)r^{2n-2}} - \frac{2\Lambda}{n(n+1)}r^2, \\ \omega_n &= \frac{16\pi}{nVol(S^n)} = \frac{8\Gamma(\frac{n+1}{2})}{n\pi^{\frac{n-1}{2}}}, \end{aligned} \tag{2.2}$$

in which Q is the electric/magnetic charge of Maxwell field, $\Lambda > 0$ -de Sitter$\Lambda < 0$ -anti-de Sitter. For general M and Q, $l = \sqrt{\frac{n(n+1)}{2\Lambda}}$ is the curvature radius of dS space, $Vol(S^n)$ denotes the volume of a unit n-sphere $d\Omega_n^2$, the equation $f(r) = 0$ may have four real roots when the parameters M, Q, l satisfy a condition[52]. Three of them are positive: the largest one is the cosmological horizon (CEH) $r = r_c$, the smallest is the inner (Cauchy) horizon of black hole, the middle one is the outer horizon (BEH) $r = r_+$ of black hole.

$$d\Omega_n^2 = d\chi_2^2 + \sin^2\chi_2^2 d\chi_3^2 + \cdots \prod_{i=2}^{n} \sin_i^2 d\chi_{n+1}^2.$$

The equations $f(r_+) = 0$ and $f(r_c) = 0$ are rearranged to

$$\begin{aligned}
Q^2 &= \frac{8(n-1)(r_+ r_c)^{n-1}}{n\omega_n^2}\left(1 - \frac{2(r_c^{n+1} - r_+^{n+1})}{n(n+1)(r_c^{n-1} - r_+^{n-1})}\Lambda\right), \\
\frac{2\Lambda}{n(n+1)} &= \left(1 - \frac{n\omega_n^2 Q^2}{8(n-1)(r_c r_+)^{n-1}}\right)\left(\frac{r_c^{n-1} - r_+^{n-1}}{r_c^{n+1} - r_+^{n+1}}\right), \qquad (2.3) \\
\omega_n M &= (r_c^{n-1} + r_+^{n-1}) - \frac{2\Lambda(r_c^{2n} - r_+^{2n})}{n(n+1)(r_c^{n-1} - r_+^{n-1})} \\
&= (r_c^{n-1} + r_+^{n-1}) - \frac{(r_c^{2n} - r_+^{2n})}{(r_c^{n+1} - r_+^{n+1})}\left(1 - \frac{n\omega_n^2 Q^2}{8(n-1)(r_c r_+)^{n-1}}\right) \\
&= \frac{(r_c r_+)^{n-1}(r_c^2 - r_+^2)}{(r_c^{n+1} - r_+^{n+1})} + \frac{(r_c^{2n} - r_+^{2n})}{(r_c^{n+1} - r_+^{n+1})}\frac{n\omega_n^2 Q^2}{8(n-1)(r_c r_+)^{n-1}}. \qquad (2.4)
\end{aligned}$$

The surface gravities at the black hole horizon and the cosmological horizon are respectively

$$\kappa_+ = \frac{1}{2}\left.\frac{df(r)}{dr}\right|_{r=r_+} = \frac{1}{2r_+}\left((n-1) - \frac{2\Lambda}{n}r_+^2 - \frac{n\omega_n^2 Q^2}{8r_+^{2n-2}}\right), \qquad (2.5)$$

$$\kappa_c = \frac{1}{2}\left.\frac{df(r)}{dr}\right|_{r=r_c} = \frac{1}{2r_c}\left((n-1) - \frac{2\Lambda}{n}r_c^2 - \frac{n\omega_n^2 Q^2}{8r_c^{2n-2}}\right). \qquad (2.6)$$

The thermodynamic quantities corresponding to the two horizons satisfy the first law of thermodynamics respectively[22, 44, 53, 54]

$$\delta M = \frac{\kappa_+}{2\pi}\delta S_+ + \Phi_+ \delta Q + V_+ \delta P, \qquad (2.7)$$

$$\delta M = \frac{\kappa_c}{2\pi}\delta S_c + \Phi_c \delta Q + V_c \delta P, \qquad (2.8)$$

where

$$S_+ = \frac{r_+^n Vol(S^n)}{4G}, \quad V_+ = \frac{Vol(S^n)}{n+1}r_+^{n+1}, \quad \Phi_+ = \frac{n}{4(n-1)}\frac{\omega_n Q}{r_+^{n-1}}, \quad S_c = \frac{r_c^n Vol(S^n)}{4G},$$

$$V_c = \frac{Vol(S^n)}{n+1}r_c^{n+1}, \quad \Phi_c = -\frac{n}{4(n-1)}\frac{\omega_n Q}{r_c^{n-1}}, \quad P = -\frac{n\Lambda}{16\pi}. \qquad (2.9)$$

III. THERMODYNAMIC QUANTITY OF RN-DS SPACETIME

In the above section, we have obtained thermodynamic quantities without considering the relationship between the black hole horizon and the cosmological horizon. Because there are three variables M, Q and Λ in the spacetime, the thermodynamic quantities corresponding to the black hole horizon and the cosmological horizon are function with respect to M, Q and Λ. The thermodynamic quantities corresponding to the black hole horizon are related to the ones corresponding to the cosmological horizon. When the thermodynamic property of charged de Sitter spacetime is studied, we must consider the relationship with the two horizon, Recently, by studying Hawking radiation of de Sitter spacetime, in [55, 56] it is obtained that the outgoing rate of the charged de Sitter spacetime which radiates particles with energy ω is

$$\Gamma = e^{\Delta S_+ + \Delta S_c}, \tag{3.1}$$

where ΔS_+ and ΔS_c are Bekenstein-Hawking entropy difference corresponding to the black hole horizon and the cosmological horizon after the charged de Sitter spacetime radiates particles with energy ω. Therefore, the thermodynamic entropy of the charged de Sitter spacetime is the sum of the black hole horizon entropy and the cosmological horizon entropy

$$S = S_+ + S_c. \tag{3.2}$$

Substitute Eqs.(2.7) and (2.8) into Eq.(3.2), we can get

$$dS = 2\pi\left(\frac{1}{\kappa_+}+\frac{1}{\kappa_c}\right)dM - \frac{n\pi\omega_n Q}{2(n-1)}\left(\frac{1}{r_+^{n-1}\kappa_+}+\frac{1}{r_c^{n-1}\kappa_c}\right)dQ + \frac{Vol(S^n)}{4(n+1)}\left(\frac{r_+^{n+1}}{\kappa_+}+\frac{r_c^{n+1}}{\kappa_c}\right)d\Lambda. \tag{3.3}$$

For simplicity, we set a fixed Q . In this case the above equation turns into

$$dS = 2\pi\left(\frac{1}{\kappa_+}+\frac{1}{\kappa_c}\right)dM + \frac{Vol(S^n)}{4(n+1)}\left(\frac{r_+^{n+1}}{\kappa_+}+\frac{r_c^{n+1}}{\kappa_c}\right)d\Lambda. \tag{3.4}$$

From Eqs.(2.3), (2.4), (2.5) and (2.6) one can obtain

$$\begin{aligned} dr_c &= \frac{1}{2\kappa_c}\left(\frac{\omega_n dM}{r_c^{n-1}} - \frac{n\omega_n^2 Q dQ}{4(n-1)r_c^{2n-2}} + \frac{2r_c^2}{n(n+1)}d\Lambda\right), \\ dr_+ &= \frac{1}{2\kappa_+}\left(\frac{\omega_n dM}{r_+^{n-1}} - \frac{n\omega_n^2 Q dQ}{4(n-1)r_+^{2n-2}} + \frac{2r_+^2}{n(n+1)}d\Lambda\right). \end{aligned} \tag{3.5}$$

Recently, in [22] the thermodynamic volume of higher dimensional RN-dS black hole is given as

$$V = \frac{Vol(S^n)}{n+1}\left(r_c^{n+1} - r_+^{n+1}\right). \tag{3.6}$$

Substitute Eq. (3.5) into Eq. (3.6), one can get

$$dV = \frac{\omega_n Vol(S^n)}{2}\left(\frac{r_c}{\kappa_c} - \frac{r_+}{\kappa_+}\right)dM + \frac{Vol(S^n)}{n(n+1)}\left(\frac{r_c^{n+2}}{\kappa_c} - \frac{r_+^{n+2}}{\kappa_+}\right)d\Lambda$$
$$- \frac{2\pi\omega_n Q}{(n-1)}\left(\frac{1}{r_c^{n-2}\kappa_c} - \frac{1}{r_+^{n-2}\kappa_+}\right)dQ. \tag{3.7}$$

Substituting Eq. (3.7) into Eq. (3.3), one can obtain the thermodynamic equation for the thermodynamic quantities of higher dimensional RN-dS black hole[45].

$$dM = T_{eff}dS - P_{eff}dV + \varphi_{eff}dQ, \tag{3.8}$$

where

$$T_{eff} = \frac{B_1}{4\pi r_c x(1+x)} + \frac{n\omega_n^2 Q^2 B_2}{32\pi r_c^{2n-1}(n-1)x^{2n-1}(1+x)}, \tag{3.9}$$

$$P_{eff} = \frac{B_3}{16\pi r_c^2 x(1+x)} + \frac{n^2\omega_n^2 Q^2 B_4}{16\times 8\pi(n-1)r_c^{2n}x^{2n-1}(1+x)}, \tag{3.10}$$

$$\varphi_{eff} = \frac{n\omega_n Q}{4(n-1)}\frac{(1-x^{2n})}{r_c^{n-1}x^{n-1}(1-x^{n+1})}, \tag{3.11}$$

and

$$(1-x^{n+1})B_1 = n(1-x^2)(1+x^{n+1}) - (1+x^2)(1-x^{n+1}),$$
$$(1-x^{n+1})B_2 = (1-x^{2n})(1+x^{n+1}) - n(1+x^{2n})(1-x^{n+1}),$$
$$(1-x^{n+1})^2 B_3 = n[n(1-x^2)(1+x^{2n+1}) - (1+x^2)(1-x^{2n+1}) + 2x^{n+1}(1-x)],$$
$$(1-x^{n+1})^2 B_4 = 2nx^{n+1}(1-x^{2n-1}) - (n-1)(1-x^{4n+1}) - (n+1)x^{2n}(1-x). \tag{3.12}$$

$x := r_+/r_c$ and $0 < x < 1$.[1]

The thermodynamic quantities defined in (3.9), (3.10) and (3.11) meet the thermodynamic equation (3.8). When $x \to 1$, namely the both horizons coincide,

$$P_{eff} \to 0,$$
$$T_{eff} \to \frac{(n-1)}{4\pi r_c(n+1)} - \frac{n^2\omega_n^2 Q^2}{32\pi r_c^{2n-1}(n+1)} = \frac{1}{4\pi r_c(n+1)}\left(2r_c^2\Lambda - (n-1)^2\right). \tag{3.13}$$

In this case, from Eqs. (2.3) and (2.4), we have

$$Q^2 = \frac{8(n-1)r_c^{2(n-1)}}{n\omega_n^2}\left(1 - \frac{2r_c^2}{n(n-1)}\Lambda\right), \quad \omega_n M = 2r_c^{n-1} - \frac{4r_c^{n+1}\Lambda}{(n+1)(n-1)}, \tag{3.14}$$

[1] It should be noted that the parameters M, Q, l should satisfy a condition to guarantee four real roots to exist. Under the condition, the x here cannot tend to zero.

Due to $Q^2 \geq 0, M \geq 0$, thus $\frac{2r_c^2\Lambda}{n(n-1)} \leq 1$. When $M^2 = Q^2$, from Eq. (3.14) one can obtain

$$2r_c^2\Lambda = \frac{(n+1)(n-1)}{n^2}\left(1+\sqrt{1+n^3(n-2)}\right). \tag{3.15}$$

Therefore, when $M^2 \geq Q^2$, there should be the relation $2r_c^2\Lambda \geq \frac{(n+1)(n-1)}{n^2}\left(1+\sqrt{1+n^3(n-2)}\right)$. In Fig.1 it shows

$$\frac{(n+1)(n-1)}{n^2}\left(1+\sqrt{1+n^3(n-2)}\right) \geq (n-1)^2. \tag{3.16}$$



FIG. 1: The plot to verify the effective temperature $T_{eff} > 0$.

According to Eq. (3.13), $T_{eff} > 0$, which fulfills the stability condition of thermodynamic system. However, the problem of considering the black hole horizon and the cosmological one as independent each other is when the two horizons coincide , namely $\kappa_+ = \kappa_c = 0$, the temperature on the black hole horizon and the cosmological horizon are both zero, but the both horizons have nonzero area, which means that the entropy for the two horizons should not be zero. The total entropy should be

$$S = S_+ + S_c = 2S_+ = 2S_c.$$

This conclusion is inconsistent with Nernst theorem. In this case the volume-thermodynamic system becomes area-thermodynamic one. According to Eq. (3.13) the pressure of thermodynamic membrane is zero, but the temperature of thermodynamic membrane is nonzero. This can partly solve the problem that extreme de Sitter black holes do not satisfy the Nernst theorem. When $n = 2$, Eqs. (3.9) and (3.10) return to the known result.

IV. PHASE TRANSITION IN CHARGED DS BLACK HOLE SPACETIME

The investigation on the phase transition of black holes have aroused many interest. In particular, the critical behaviors of some black holes in asymptotically AdS space are similar to ones of the Van der Waals equation[16–19, 23–26]. For black holes in dS space there exist more than one horizons and the multiple horizons correspond to different thermodynamic systems. Thus they are generally non-equilibrium systems. The research on the phase transition of this kind of non-equilibrium system is few. Based on the above section, we analyze the phase transition of higher dimensional RN-dS black hole. Firstly, we compare the effective thermodynamic quantities of higher dimensional RN-dS black hole with the der Waals equation. Secondly, we will discuss the critical behaviors of thermodynamic quantities of the RN-dS black hole at constant temperature. At last, the nature of the phase transition by Ehrenfest's equations will be analyzed.

Comparing with the Van der Waals equation

$$\left(P+\frac{a}{v^2}\right)(v-b)=kT, \tag{4.1}$$

here, $v=V/N$ is the specific volume of the fluid, P its pressure, T its temperature, and k is the Boltzmann constant. From Eq.(4.1)one can depict the $P-v$ curves for fixed T. Employing the conditions and the equations the critical points satisfied, one can derive the critical temperature, the critical pressure and the critical volume. To compare with the Van der Waals equation,we set $P_{eff}\to P$ and discuss the phase transition and critical phenomena when Q is invariant.

Substituting Eq.(3.9) into Eq.(3.10), we have

$$P_{eff}=T_{eff}\frac{B_4}{2r_cB_2}+\frac{B_2B_3-B_1B_4}{8\pi r_c^2x(1+x)B_2}, \tag{4.2}$$

in which x is a dimensionless parameter. We take the specific volume of the higher dimensional RN-dS spacetime as

$$v=r_c(1-x). \tag{4.3}$$

According to

$$\left(\frac{\partial P_{eff}}{\partial v}\right)_{T_{eff}}=0,\quad \left(\frac{\partial^2 P_{eff}}{\partial v^2}\right)_{T_{eff}}=0, \tag{4.4}$$

we can calculate the position of the critical points.

Here we take $Q = 1,\ 3,\ 10$ respectively and calculate the various quantities at the critical points in different spacetime dimensions ($n = 3,\ 4,\ 5$). The results are shown in Table I.

	$n=3$			$n=4$			$n=5$		
	$Q=1$	$Q=3$	$Q=10$	$Q=1$	$Q=3$	$Q=10$	$Q=1$	$Q=3$	$Q=10$
x^c	0.769263	0.769263	0.769263	0.798722	0.798722	0.798722	0.821785	0.821785	0.821785
r_c^c	1.61337	2.79444	5.10192	1.31405	1.89519	2.83103	1.21688	1.60151	2.16396
T_{eff}^c	0.030460	0.017586	0.009232	0.047178	0.032711	0.021898	0.057924	0.044013	0.032573
P_{eff}^c	0.008375	0.002791	0.000837	0.022521	0.010827	0.004852	0.038737	0.022365	0.012249
v^c	0.372263	0.644779	1.1772	0.264489	0.381459	0.569823	0.216866	0.285412	0.385648
V^c	21.7266	195.54	2172.66	13.9192	86.8592	646.071	11.6117	60.336	367.193
M^c	1.26609	3.79828	12.6609	1.38401	4.15202	13.8401	1.51006	4.5319	15.1006
S^c	30.1578	156.705	953.675	27.6023	119.428	594.674	28.436	112.272	505.672
G^c	0.52945	1.58835	5.2945	0.395263	1.18579	3.95263	0.312726	0.938178	3.12726

TABLE I: Numerical solutions for x^c, r_c^c, T_{eff}^c, P_{eff}^c, v^c, V^c, M^c, S^c and G^c for given values of $Q = 1, 3, 10$ and spacetime dimension $n = 3, 4, 5$ respectively.

According to the above calculations, the position x^c of the critical point of the higher dimensional RN-dS black holes is independent of the electric charge. However, it is relevant to the dimension of spacetime and increases with the dimension. The position of the critical cosmological horizon r_c^c increases with the electric charge Q for fixed spacetime dimension, and decrease with the increase of the spacetime dimension for fixed electric charge. Both the critical effective temperature T_{eff}^c and the critical effective pressure P_{eff}^c decrease with the increase of the electric charge Q for fixed spacetime dimension, and increase with the spacetime dimension for fixed electric charge. The critical specific volume v^c increase with the increase of the electric charge Q for fixed spacetime dimension, and decrease with the spacetime dimension for fixed electric charge.

In order to describe the relation of P_{eff} and v in the vicinity of critical temperature, we plot the curves of P_{eff}-v at different temperatures. For the sake of brevity, we only depict the curves for the $n = 5$ case. In the cases with $n = 3,\ 4$ the behaviors are similar.

From Figs.2, when the effective temperature $T_{eff} > T_{eff}^c$, the stable condition $\left(\frac{\partial P}{\partial v}\right)_{T_{eff}} <$



(a) $Q = 1$ (b) $Q = 3$ (c) $Q = 10$

FIG. 2: *The $P_{eff} - v$ curves for $Q = 1, 3, 10$ respectively. From top to the bottom the curves correspond to the effective temperature $T^c_{eff} + 0.02$, $T^c_{eff} + 0.01$, T^c_{eff}, $T^c_{eff} - 0.01$ and $T^c_{eff} - 0.02$.*

0 can be satisfied. In the case of $T_{eff} < T^c_{eff}$, only when the value of v is large, the stable condition $\left(\frac{\partial P}{\partial v}\right)_{T_{eff}} < 0$ can be satisfied. When the system lies at a small v, $\left(\frac{\partial P}{\partial v}\right)_{T_{eff}} > 0$, thus the system is unstable. So phase transition may occur only at $T_{eff} = T^c_{eff}$.

V. ANALYTICAL CHECK OF THE CLASSICAL EHRENFEST EQUATIONS IN THE EXTENDED PHASE SPACE

According to Ehrenfest's classification, when the chemical potential and its first derivative are continuous, whereas the second derivative of chemical potential is discontinuous, this kind of phase transition is called the second-order phase transition. For Van der Waals system there is no latent heat and the liquid-gas structure do not change suddenly at the critical point. Therefor this kind of phase transition belongs to the second-order phase transition and continuous phase transition . Below we will discuss the behaviors of higher dimensional RN-dS system near the phase transition point.

We can calculate the specific heat of RN-dS system at constant pressure C_P, the volume expansivity β, and the isothermal compressibility κ

$$C_P = T_{eff}\left(\frac{\partial S}{\partial T_{eff}}\right)_{P_{eff}} = -T_{eff}\frac{\partial^2 \mathrm{G}}{\partial T^2_{eff}}$$

$$= \frac{Vol(S^n) r_c^{n-1}}{4} n T_{eff} \left(\frac{r_c(1+x^{n-1})\left(\frac{\partial P_{eff}}{\partial r_c}\right)_x - (1+x^n)\left(\frac{\partial P_{eff}}{\partial x}\right)_{r_c}}{\left(\frac{\partial T_{eff}}{\partial x}\right)_{r_c}\left(\frac{\partial P_{eff}}{\partial r_c}\right)_x - \left(\frac{\partial T_{eff}}{\partial r_c}\right)_x \left(\frac{\partial P_{eff}}{\partial x}\right)_{r_c}} \right), \qquad (5.1)$$

$$\beta = \frac{1}{v}\left(\frac{\partial v}{\partial T_{eff}}\right)_{P_{eff}} = \frac{1}{v}\frac{\partial^2 \mu}{\partial T_{eff}\partial P_{eff}}$$

$$= -\frac{1}{v}\left(\frac{r_c\left(\frac{\partial P_{eff}}{\partial r_c}\right)_x + (1-x)\left(\frac{\partial P_{eff}}{\partial x}\right)_{r_c}}{\left(\frac{\partial T_{eff}}{\partial x}\right)_{r_c}\left(\frac{\partial P_{eff}}{\partial r_c}\right)_x - \left(\frac{\partial T_{eff}}{\partial r_c}\right)_x\left(\frac{\partial P_{eff}}{\partial x}\right)_{r_c}}\right), \tag{5.2}$$

$$\kappa = -\frac{1}{v}\left(\frac{\partial v}{\partial P_{eff}}\right)_{T_{eff}} = -\frac{1}{v}\frac{\partial^2 \mu}{\partial P_{eff}^2}$$

$$= -\frac{1}{v}\left(\frac{r_c\left(\frac{\partial T_{eff}}{\partial r_c}\right)_x + (1-x)\left(\frac{\partial T_{eff}}{\partial x}\right)_{r_c}}{\left(\frac{\partial P_{eff}}{\partial r_c}\right)_x\left(\frac{\partial T_{eff}}{\partial x}\right)_{r_c} - \left(\frac{\partial P_{eff}}{\partial x}\right)_{r_c}\left(\frac{\partial T_{eff}}{\partial r_c}\right)_x}\right), \tag{5.3}$$

where $S = \frac{Vol(S^n)}{4} r_c^n (1 + x^n)$.

We also depict the curves of $C_P - x$, $\beta - x$ and $\kappa - x$ in the FIG 3, FIG 4 and FIG 5 respectively. From these curves, we find that the specific heat of higher dimensional RN-dS system at constant pressure C_P, the expansion coefficient β, and the compressibility κ exist infinite peak. While the curves of $S - x$ and $G - x$ in FIG 6 and FIG 7 show that the Gibbs function G and the entropy S are both continuous at the critical point. According to Ehrenfest, the phase transition of the higher dimensional RN-dS black hole should be the second-order one, which is similar to the four dimensional case.



(a) $Q = 1$ (b) $Q = 3$ (c) $Q = 10$

FIG. 3: *$C_P - x$ curves for RN-dS black hole with $n = 5$ corresponding to the critical effective pressure $P^c_{eff} = 0.038737$, $P^c_{eff} = 0.022365$ and $P^c_{eff} = 0.012249$ respectively.*

Ehrenfest's equations (slope formula) for RN-dS black holes are

$$\left(\frac{\partial P_{eff}}{\partial T_{eff}}\right)_S = \frac{C_P^2 - C_P^1}{v^c T^c_{eff}(\beta_2 - \beta_1)} = \frac{\Delta C_P}{v^c T^c_{eff}\Delta\beta}, \tag{5.4}$$



(a) $Q = 1$ (b) $Q = 3$ (c) $Q = 10$

FIG. 4: *$\beta - x$ curves for RN-dS black hole with $n = 5$ corresponding to the critical effective pressure $P^c_{eff} = 0.038737$, $P^c_{eff} = 0.022365$ and $P^c_{eff} = 0.012249$ respectively.*



(a) $Q = 1$ (b) $Q = 3$ (c) $Q = 10$

FIG. 5: *$\kappa - x$ curves for RN-dS black hole with $n = 5$ corresponding to the critical effective temperature $T^c_{eff} = 0.057924$, $T^c_{eff} = 0.044013$ and $T^c_{eff} = 0.032573$ respectively.*

$$\left(\frac{\partial P_{eff}}{\partial T_{eff}}\right)_v = \frac{\beta_2 - \beta_1}{\kappa_T^2 - \kappa_T^1} = \frac{\Delta\beta}{\Delta\kappa_T}, \tag{5.5}$$

in which the subscript 1 and 2 represent phase 1 and 2 respectively.

From Maxwell relation

$$\left(\frac{\partial v}{\partial S}\right)_P = \left(\frac{\partial T}{\partial P}\right)_S \tag{5.6}$$

and Eqs. (5.4), we have

$$\left[\left(\frac{\partial P_{eff}}{\partial T_{eff}}\right)_S\right]^c = \left[\left(\frac{\partial S}{\partial v}\right)_{P_{eff}}\right]^c, \tag{5.7}$$



(a) $Q = 1$ (b) $Q = 3$ (c) $Q = 10$

FIG. 6: *$S-T$ curves for RN-dS black hole with $n = 5$ corresponding to the critical effective pressure $P^c_{eff} = 0.038737$, $P^c_{eff} = 0.022365$ and $P^c_{eff} = 0.012249$ respectively.*



(a) $Q = 1$ (b) $Q = 3$ (c) $Q = 10$

FIG. 7: *$G-T$ curves for RN-dS black hole with $n = 5$ corresponding to the critical effective pressure $P^c_{eff} = 0.038737$, $P^c_{eff} = 0.022365$ and $P^c_{eff} = 0.012249$ respectively. Here the Gibbs free energy $G = M - T_{eff}S - \varphi_{eff}Q$.*

thus

$$\frac{\Delta C_P}{T^c_{eff} v^c \Delta\beta} = \left[\left(\frac{\partial S}{\partial v}\right)_{P_{eff}}\right]^c . \tag{5.8}$$

Note that the footnote "c" denotes the values of physical quantities at critical point in our letter. Here

$$\left(\frac{\partial S}{\partial v}\right)_{P_{eff}} = \left(\frac{\left(\frac{\partial S}{\partial x}\right)_{r_c}\left(\frac{\partial P_{eff}}{\partial r_c}\right)_x - \left(\frac{\partial S}{\partial r_c}\right)_x\left(\frac{\partial P_{eff}}{\partial x}\right)_{r_c}}{\left(\frac{\partial v}{\partial x}\right)_{r_c}\left(\frac{\partial P_{eff}}{\partial r_c}\right)_x - \left(\frac{\partial v}{\partial r_c}\right)_x\left(\frac{\partial P_{eff}}{\partial x}\right)_{r_c}}\right) . \tag{5.9}$$

From Maxwell relation

$$\left(\frac{\partial v}{\partial S}\right)_T = \left(\frac{\partial T}{\partial P}\right)_v \tag{5.10}$$

, and Eqs. (5.5), one can get

$$\frac{\Delta\beta}{\Delta\kappa_T} = \left(\frac{\partial v}{\partial S}\right)^c_{T_{eff}}, \tag{5.11}$$

here

$$\left(\frac{\partial S}{\partial v}\right)_{T_{eff}} = \left(\frac{\left(\frac{\partial S}{\partial x}\right)_{r_c}\left(\frac{\partial T_{eff}}{\partial r_c}\right)_x - \left(\frac{\partial S}{\partial r_c}\right)_x\left(\frac{\partial T_{eff}}{\partial x}\right)_{r_c}}{\left(\frac{\partial v}{\partial x}\right)_{r_c}\left(\frac{\partial T_{eff}}{\partial r_c}\right)_x - \left(\frac{\partial v}{\partial r_c}\right)_x\left(\frac{\partial T_{eff}}{\partial x}\right)_{r_c}}\right). \tag{5.12}$$

According to Eq. (5.2), when $r_c\left(\frac{\partial T_{eff}}{\partial r_c}\right)_x + (1-x)\left(\frac{\partial T_{eff}}{\partial x}\right)_{r_c} \neq 0$, the critical points satisfy

$$\left(\frac{\partial P_{eff}}{\partial r_c}\right)_x\left(\frac{\partial T_{eff}}{\partial x}\right)_{r_c} - \left(\frac{\partial P_{eff}}{\partial x}\right)_{r_c}\left(\frac{\partial T_{eff}}{\partial r_c}\right)_x = 0. \tag{5.13}$$

Substituting Eq. (5.13) into Eq. (5.12) and comparing with Eq. (5.9), we have

$$\left(\frac{\partial S}{\partial v}\right)^c_{T_{eff}} = \left(\frac{\partial S}{\partial v}\right)^c_{P_{eff}}. \tag{5.14}$$

So far, we have proved that both the Ehrenfest equations are correct at the critical point. Utilizing Eq .(5.14), the Prigogine–Defay (PD) ratio (Π)can be calculated as

$$\Pi = \frac{\Delta C_p \Delta\kappa_T}{T^c_{eff} v^c (\Delta\beta)^2} = 1. \tag{5.15}$$

Hence the phase transition occurring at $T_{eff} = T^c_{eff}$ is a second order equilibrium transition. This is true in spite of the fact that the phase transition curves are smeared and divergent near the critical point. This result is in agreement with the one in[16, 35, 37–40] about AdS black holes.

VI. CONCLUSIONS

After introducing the connection between the thermodynamic quantities corresponding to the black hole horizon and the cosmological horizon, we give the effective thermodynamic quantities of the higher dimensional RN-dS system, (3.9),(3.10) and (3.11). When describing the higher dimensional RN-dS system by the effective thermodynamic quantities, it exhibits a similar phase transition to Van der Waals equation. In Sec.4 it is shown that the position x of the phase transition point in the higher dimensional RN-dS system is irrelevant to the

electric charge of the system. This indicates that for fixed charge when the ratio of the black hole horizon and the cosmological horizon is x^c, the second-order phase transition will occur. There are some differences between the higher dimensional RN-dS black hole and the charged AdS black hole. According to the isothermal curves of the Van der Waals equation and charged AdS black holes when the temperature is lower than the critical one there exist two different phases for some values of the volume, namely coexistence region. From Figs.2, for the higher dimensional RN-dS black hole no two-phase region exists for any temperature. When the effective temperature $T_{eff} > T^c_{eff}$, $\left(\frac{\partial P_{eff}}{\partial v}\right)_{T_{eff}} < 0$ which satisfy the stable condition. When $T_{eff} < T^c_{eff}$, at some intervals the isothermal curve corresponds to $\left(\frac{\partial P_{eff}}{\partial v}\right)_{T_{eff}} > 0$. At this time the system is unstable. The states with $\left(\frac{\partial P_{eff}}{\partial v}\right)_{T_{eff}} > 0$ turn up at small value of v with $v = r_c(1-x)$, namely at some interval of x with larger values. This means that when the both horizons approach, the system lies at a non-equilibrium state. Therefore, for the higher dimensional RN-dS black hole the state with coincided horizons does not exist.

In Sec. 5 we analyzed the phase transition of higher dimensional RN-dS system. It shows that at the critical point the specific heat at constant pressure, the volume expansivity β and the isothermal compressibility κ of the RN-dS system exist infinite peak, while the entropy and the Gibbs potential G are continuous. Therefore for the phase transition of the RN-dS system no latent heat and no specific volume changes suddenly, it belongs to the second-order phase transition.

We carry out an analytical check of Ehrenfest equations and prove that both Ehrenfest equations are satisfied. This result is consistent with the nature of liquid–gas phase transition at the critical point, hence deepening the understanding of the analogy of RN-dS spacetime and liquid-gas systems.

Acknowledgments

This work is supported by NSFC under Grant Nos.(11175109;11075098;11247261;11205097) and the doctoral Sustentation foundation of Shanxi Datong University (2011-B-03).

[1] J. D. Bekenstein, Lett. Nuovo Cimento 4, 737 (1972).

[2] J. D. Bekenstein, Phys. Rev. D 7, 949 (1973).

[3] J. D. Bekenstein, Phys. Rev. D 9, 3292 (1974).

[4] J. M. Bardeen, B. Carter, S. W. Hawking, Commun. Math. Phys. 31, 161 (1973).

[5] S. W. Hawking, Nature, 248, 30 (1974).

[6] S. W. Hawking, Commun. Math. Phys. 43 199 (1975).

[7] S. Hawking, D. N. Page, Commun. Math. Phys. 87, 577 (1983).

[8] A. Chamblin, R. Emparan, C. Johnson, and R. Myers, Phys.Rev. D 60, 064018 (1999), [arXiv:hep-th/9902170].

[9] A. Chamblin, R. Emparan, C. Johnson, and R. Myers, Phys.Rev. D 60,104026 (1999) , [arXiv:hep-th/9904197].

[10] O. Aharony, S. S. Gubser, J. M. Maldacena, H. Ooguri, and Y. Oz, Phys. Rept. 323, 183 (2000), [arXiv:hep-th/9905111].

[11] S. S. Gubser, Phys. Rev. D 78, 065034 (2008), [arXiv:0801.2977[hep-th]].

[12] S. A. Hartnoll, C. P. Herzog, and G. T. Horowitz, Phys. Rev. Lett. 101, 031601 (2008), [arXiv:0803.3295[hep-th]].

[13] A. Sahay, T. Sarkar, and G. Sengupta, JHEP 1004, 118 (2010), [arXiv:1002.2538[hep-th]].

[14] A. Sahay, T. Sarkar, and G. Sengupta, JHEP 1007, 082 (2010), [arXiv:1004.1625[hep-th]].

[15] A. Sahay, T. Sarkar, and G. Sengupta, JHEP 1011, 125 (2010), [arXiv:1009.2236[hep-th]].

[16] R. Banerjee, S. Ghosh, and D. Roychowdhury, Phys. Lett. B 696, 156 (2011), [arXiv:1008.2644[hep-th]].

[17] D. Kastor, S. Ray, J. Traschen, Class. Quant. Grav. 26, 95011 (2009), [arXiv:0904.2765[hep-th]]

[18] B. P. Dolan, Class. Quant. Grav. 28, 125020 (2011), [arXiv:1008.5023[hep-th]].

[19] S. Gunasekaran, D. Kubiznak, R. B. Mann, JHEP, 1211, 110 (2012), [arXiv:1208.6251.[hep-th]].

[20] B. P. Dolan, Phys. Rev. D 84, 127503 (2011), [arXiv:1109.0198.[hep-th]].

[21] M. Cvetic, G.W. Gibbons, D. Kubiznak, C. N. Pope, Phys. Rev. D 84, 024037 (2011), [arXiv:1012.2888[hep-th]].

[22] B. P. Dolan, D Kastor, D. Kubiznak, R. B. Mann, J. Traschen, Phys. Rev. D 87, 104017 (2013), [arXiv:1301.5926[hep-th]].

[23] N. Altamirano, D. Kubiznak, R. B. Mann and Z. Sherkatghanad, arXiv:1401.2586 [hep-th].

[24] B. P. Dolan, arXiv:1209.1272[gr-qc].

[25] D. C. Zou, S. J. Zhang, B. Wang, Phys. Rev. D 89, 044002 (2014), [arXiv:1311.7299[hep-th]].

[26] D. Kubiznak, R. B. Mann, JHEP 1207, 033 (2012), [arXiv:1205.0559[hep-th]].

[27] S. W. Wei, Y. X. Liu, Phys. Rev. D 87, 044014 (2013), [arXiv:1209.1707[gr-qc]].

[28] R. G. Cai, L. M. Cao, L. Li, and R. Q. Yang, JHEP, 1309, 005 (2013), [arXiv:1306.6233[gr-qc]].

[29] Shao-Wen Wei, Yu-Xiao Liu, [arXiv:1402.2837[gr-qc]].

[30] S. H. Hendi, M. H. Vahidinia, Phys. Rev. D 88, 084045 (2013), [arXiv:1212.6128[hep-th]].

[31] E. Spallucci, A. Smailagic, Phys. Lett. B 723, 436 (2013), [arXiv:1305.3379[hep-th]].

[32] A. Belhaj, M. Chabab, H. E. Moumni and M. B. Sedra, arXiv:1306.2518[hep-th].

[33] R. Zhao, H. H. Zhao, M. S. Ma and L. C. Zhang, Eur. Phys. J. C 73, 2645 (2013), [arXiv:1305.3725[gr-qc]].

[34] M. B. J. Poshteh, B. Mirza, Z.Sherkatghanad, Phys. Rev. D 88, 024005 (2013),[arXiv:1306.4516[gr-qc]].

[35] J.-X. Mo, W.-B. Liu, Phys. Lett. B 727,336 (2013).

[36] J.-X. Mo, W.-B. Liu, [arXiv:1401.0785[gr-qc]].

[37] R. Banerjee, S. K. Modak, S. Samanta, Phys. Rev. D84, 064024 (2011), [arXiv:1005.4832[hep-th]].

[38] R. Banerjee, D. Roychowdhury, JHEP 11, 004 (2011), [arXiv:1109.2433[hep-th]].

[39] R. Banerjee, D. Roychowdhury, Phys. Rev. D 85, 044040 (2012), [arXiv:1111.0147[hep-th]].

[40] R. Banerjee, D. Roychowdhury, Phys. Rev. D 85, 104043 (2012), [arXiv:1203.0118[hep-th]].

[41] N. Altamirano, D. Kubiznak and R. Mann, Phys. Rev. D 88, 101502 (2013), [arXiv:1306.5756 [hep-th]].

[42] M. S. Ma, H. H. Zhao, L. C. Zhang, R. Zhao, Int. J. Mod. Phys. **A**, to be published.

[43] R. G. Cai, Nucl. Phys. **B**628, 375 (2002).

[44] Y. Sekiwa, Phys. Rev. D 73, 084009(2006),[arXiv:hep-th/0602269].

[45] M. Urano, A. Tomimatsu, Class. Quant. Grav.26, 105010 (2009), [arXiv:0903.4230[hep-th]].

[46] L. C. Zhang, H. F. Li and R. Zhao, Science China, Physics, Mechanics Astronomy 54, 1384 (2011).

[47] Y. S. Myung, Phys. Rev. D 77, 104007 (2008).

[48] M. Eune, W. Kim, Phys. Lett. B 723, 177 (2013).

[49] R. G. Cai, Physics 34, 555(2005),(in Chinese).

[50] S. Bhattacharya, A. Lahir, Eur. Phys. J. C 73, 2673 (2013), [arXiv:1301.4532[gr-qc]].

[51] R. G. Cai, J. Y. Ji, K. S. Soh, Classical. Quantum. Grav. 15, 2783 (1998).

[52] B. D. Koberlein and R. L. Mallett, Phys. Rev. D 49, 5111(1994).

[53] G. W. Gibbons, H. Lü , D. N. Page, and C. N. Pope, Phys. Rev. Lett. 93, 171102 (2004), [arXiv:hep-th/0409155].

[54] G. W. Gibbons, H. Lü , D. N. Page, and C. N. Pope, J. Geom. Phys. 53, 49(2005), [arXiv:hep-th/0404008].

[55] R. Zhao, L. C. Zhang, H. F. Li, Gen. Relat. Grav. 42, 975 (2010).

[56] R. Zhao, L. C. Zhang, H. F. Li and Y. Q. Wu, Eur. Phys. J. C. 65, 289 (2010).